



ISESER 2020

**INTERNATIONAL SYMPOSIUM
FOR ENVIRONMENTAL SCIENCE
AND ENGINEERING RESEARCH:**

ISESER2020

PROCEEDING

ISBN 978-605-83522-2-3



04-05 July 2020, Manisa, Turkey

O 15. INVESTIGATION OF THE EFFECT OF CORONA VIRUS OUTBREAK MEASURES ON ATMOSPHERIC PM VALUES FOR KONYA CITY CENTRE

Alaa Motar Shayia^{1*}, Hüseyin Toros², Mustafa Sağdıç³, Şükrü Dursun⁴

¹ *Atmospheric Sciences Dept., College of Science, Mustansiriyah University, Baghdad, Iraq,
+9647704671275*

² *Istanbul Technical University, Faculty of Aeronautics and Astronautics, Department of Meteorology,
Maslak, 34469, Istanbul, Turkey*

³ *Yıldız Technical University, Faculty of Education, Department of Social Sciences and Turkish
Education, Istanbul, Turkey*

⁴ *Konya Technical University, Engineering and Natural Science Faculty, Environmental Engineering
Department, 42003 Konya, Turkey*

*E-mail: al.shayia.atmsc@uomustansiriyah.edu.iq, toros@itu.edu.tr, msagdic@yildiz.edu.tr,
sdursun@ktun.edu.tr*

ABSTRACT: This present study is to investigate the change in atmospheric particulate matter values of curfews taken due to the corona virus outbreak in Konya city centre. In this study, PM10 pollutant data was used in Konya city centre data of the Ministry of Environment and Urbanization. The period in which strict measures were taken in the study was compared with the period between 16 March 2020 and 15 April 2020 before and after. In addition, seasonal conditions, 2020 epidemic period data and normal period 2018 and 2019 data were also compared. As a result, because the vast majority of living in major cities in the epidemic period of restrictions in Turkey Konya seen the positive effects on air quality in the city centre it has been found to be examined in PM10. It was revealed that the measures taken in the period of virus outbreak, curtailment of traffic and reduction of industrial activities caused a remarkable improvement in air quality. Considering the measures and improvements in air quality, it is considered to be worth investigating how the measures should be evaluated in the coming periods in combating air pollution.

Keywords: *Air pollution, Konya, Corona Virus, Pandemic, Air quality*

1. INTRODUCTION

Coronavirus, a novel infectious disease, was first identified in the Wuhan province of China in December 2019 (Kanniah *et al.*, 2020; Huang *et al.*, 2020; Chen *et al.*, 2020). This disease later spread to other countries in Asia, Europe (mainly Italy, Spain, France and the United Kingdom), Africa and America (mainly the United States), and became a pandemic. COVID-19 is extremely communicable to more than 11.8 million People (confirmed cases on 09 July 2020) have been contaminated in 210 countries with more than 534, 902 recorded deaths (09 July 2020; URL-1). When countries entered the lockdown, manufacturing operations were shut down worldwide (Muhammad & Xingle, 2020). Transport is, among many other industries, the business most seriously impacted by the lockdown. Road and air transport comes to a stop because people were not permitted or hesitant to fly. According to the survey, air traffic decreased by 96 per cent attributable to COVID-19, the lowest in 75 years (CNN, 2020). As a potential side effect of this extraordinary lockdown, several countries have undergone a drastic decrease in air quality. In China, the Finnish Centre for Energy and Clean Air Study announced that steps to curb the spread of COVID-19, such as travel restrictions and plant closures, culminated in a 25% decrease in CO₂ emission (Carbon Brief 2020, URL-2) Similarly, satellite images by the European Space Agency (ESA) revealed a substantial reduction in NO₂ pollution in northern Italy between 01.January and 11.March 2020. Lockdowns to protect off coronavirus (Bao & Zhang, 2020). In addition, the Institute of environmental Science and Meteorology (IESM) has reported that since the launch of the Luzon strengthened environmental quarantine on 16 March 2020, the PM_{2.5} and PM₁₀ concentrations in Metro Manila have been substantially decreased due to reduced use in crushing and grinding machines and low road dust exposure (Wang *et al.*, 2020).

Particle matter (PM) air pollution is more complicated, spanning a broad variety of applications. This consists of a multi-component matrix derived from different anthropogenic (power production, traffic-related, etc.) and natural causes (biomass combustion, pollen, etc.) and is subject to a variety of atmospheric processes (Zoran *et al.*, 2020). In certain urban coagulated regions, PM concentrations are

typically dominated by various size fractions (ultrafine PM_{0.1} particles with a diameter of < 0.1 μm; small PM_{2.5} particles with a diameter of ≤0.2.5 μm; coarse PM₁₀ particles with a diameter of > 0.2.5 μm and ≤10 μm) (Zoran et al., 2019; Khan et al., 2019).many studied in different countries were done to investigate the impact of COVID19 on air pollution concentrations and these studies have shown that the lockdown contributed to a substantial decrease in aerosols optical depth AOD over sea and emissions over oceanic regions, although a large decrease in tropospheric NO₂ was reported over areas not influenced by seasonal biomass combustion. PM₁₀, PM_{2.5}, NO₂, SO₂ and CO concentrations showing a notable decreased in urban areas during the lock-down process relative to the same times in 2018 and 2019 (Dutheil et al., 2020; Muhammad & Xingle, 2020; Bao & Zhang, 2020; Otmani et al., 2020). The present study aims to investigate the change of PM₁₀ concentration due to the effects of coronavirus outbreak in Konya city center, Turkey using the data of the existing air quality monitoring station in the city before and after Covid19 measures.

2. MATERIAL AND METHOD

Study area and data sources

Konya is a closed basin located in the central part of the interior of Anatolia, Turkey (Figure 1). The closed basin consists of wide ovals and plateaus. As the Taurus Mountains cover the south of the basin, it prevents the moist air of the Mediterranean from coming to the region. Therefore, although it is close to the Mediterranean, it has a very arid climate and has a semi-arid feature. With this feature, summers are hot and dry and winters are cold. Turkey is much less rainfall than in the general. In the past, it called Turkey's granary and socio-cultural development, as this feature is also changed by the change of climate and water resources. Industrial areas, especially provincial center, have also developed. Since air circulation is not enough in the city center from time to time, air pollution events reach levels that disturb people on some days.

Climate: In Konya, located in the southern part of Central Anatolia, winters are harsh, cold and snowy, summers are hot and dry. The average annual temperature is 11.6 °C. The daily maximum annual temperature is 17.9 °C and the minimum annual temperature is 5.4 °C. The highest temperature is 40.6 °C and the lowest is -28.2 °C (Figure 1).




Figure 1. long term, between 1929 to 2019, average values of Temperature and precipitation of Konya city (URL-3).



Figure 1. Location of research area Konya (URL-4)

Air pollution: Depending on the development of the industry in the city center, fossil fuel consumption for energy needs is one of the emission sources thrown into the air. Again, because the region is very cold in the winter months, fossil fuels used for heating are another important source of air pollution. It is thought that it contributes to air pollution from industrial processes. Vehicles in traffic have a significant impact on air pollution, and their contribution to air pollution especially increases in some periods and hours. The exhaust gases of the vehicles are especially important in terms of very small PM, CO, and NO_x emissions.

Data: The air quality monitoring stations located in the center of Konya as an all other Turkish cities. In this study, air pollution measurement values of 4 stations in the city center of Konya were obtained from the Ministry of Environment and Urbanization WEB page (URL-5). PM₁₀, SO₂, PM_{2.5}, NO₂ and O₃ values are measured at the sampling stations of the Ministry and Konya municipality are published on the WEB page as hourly and then daily averages measurement values are given in $\mu\text{g}/\text{m}^3$.

This present study is to investigate air quality with the change in atmospheric particulate matter values of curfews taken due to the coronavirus outbreak in Konya city center. The 2nd period which is strict measures were taken for Covid-19 which were compared two periods 1 January 2020 - 15 March 2020 and 16 March 2020 - 15 April 2020. In addition, seasonal conditions, 2020 epidemic period data, and normal period 2018 and 2019 data were compared. Period 1st, before the Covid-19 measures (between January 1 and March 15). Period 2nd, after the Covid-19 struggle started (between March 16 and April 15).

R-statistic program, which is an open code statistics program, was used to create the distribution of pollution maps of the data, daily air pollutants data of 01.01.2018-15.06.2020 compared.

3. RESULTS AND DISCUSSION

Any region in the world where the industry is highly developed, provincial centers where traffic is very heavy, and settlements in the places where winter seasons are cold and using low-quality fossil fuel consumption affects the quality of human life, especially in people with respiratory diseases problems. The topographic structure and climatic features of a region are also as effective as emission sources in the formation of air pollution. Especially in the winter months, there is an increase in the emission levels as the temperature decreases with increasing fuel usage for heating systems. However, the negativity caused by meteorological conditions causes lower than expected levels of air quality problems.

Using the data of the existing air quality monitoring stations in the Konya city center: it is seen that the air quality increases and then returns to normal levels during the period when a curfew is restricted for measures taken due to the Covid-19 pandemic outbreak period. outbreak restrictions that cause the

reduction in vehicle exhaust emissions, which are important factors in the formation of some air pollutant parameters, are thought to be effective in improving the air quality in city centers.

Figure 2 shows the mean daily particle matter concentration for 4 different sampling stations in Konya city center from the beginning of 2020 to the middle of June 2020. Values in graph show a significant decreasing after pandemic outbreak middle of March 2020 except Selcuklu sampling point differentiation due to the effect of Saharan dust effect period.




Figure 2. PM₁₀ values for four different sampling stations at Meram, Karatay, Selcuklu districts of Konya city center.

When it is seen separately in Figure 3, daily mean of PM₁₀ levels as an air quality parameter in Meram district sampling station of Konya city. They show that PM₁₀ values decreased and values significantly changed for Meram regions during the outbreak of virus pandemic period.




Figure 3. PM₁₀ values for sampling stations of Meram district of Konya city center for the beginning of January 2020 to middle of June 2020.

According to Figure 4, PM₁₀ values for sampling stations in Karatay district of Konya city center are higher than Meram district but values are parallel trend with Meram region. This district includes places that are a more industrialized factory region. Values were significantly decreased during the restriction of COVID-19 pandemic after March 16th, 2020.




Figure 4. PM₁₀ values for sampling stations of Karatay district of Konya city center for the beginning of January 2020 to middle of June 2020.

When it is seen in Figure 5, PM₁₀ values for sampling stations of Selcuklu municipality region show a similar sampling period with Meram district, values of PM₁₀ and trends are similar. Improvement of air quality was seen as a similar period of a pandemic outbreak.




Figure 5. PM₁₀ values for sampling stations of Selcuklu municipality region of Konya city center for the beginning of January 2020 to middle of June 2020.

Figure 5 is shows PM₁₀ values for another sampling station of Selcuklu district in Konya city center during the period beginning of January 2020 to middle of June 2020. This region is mostly residential area including a part of a small industrial area. Most of the pollution comes from fossil fuels burning for production of energy for heating systems in homes. A small amount of pollution was introduced by industry and traffic. During cold wintertime, PM₁₀ values were increased. PM₁₀ values were significantly decreased during pandemic outbreak time and an increase was detected with the situation created by the effect of field dust and then decreased again.




Figure 6. PM₁₀ values for sampling station of Selcuklu district of Konya city center for the beginning of January 2020 to middle of June 2020.

Figure 7 is summarizing the daily mean values of PM₁₀ as an air quality parameter for Selcuklu, Konya sampling station from beginning January of 2018 to the middle of June 2020. Values showing that PM₁₀ values during summer hot period lower than colder winter periods the highest values were December 2019. Values significantly decreased during the pandemic outbreak period and after the returned to normal levels. Four month values from the middle of December 2018 are missing for some problem of the sampling equipment in the station.




Figure 7. Daily mean values of PM₁₀, as an air quality parameter of Selcuklu, Konya sampling station from beginning January of 2018 to middle of June 2020.

Daily mean values of PM₁₀, as an air quality parameter of Meram, Konya sampling station from beginning January of 2018 to middle of June 2020 (Figure 8) which shows that PM₁₀ values decreased during a pandemic outbreak. PM₁₀ values during summer hot period lower than colder winter periods and highest values were seen in December 2019.




Figure 8. Daily mean values of PM₁₀, as an air quality parameter of Meram, Konya sampling station from beginning January of 2018 to middle of June 2020.

Daily mean values of PM₁₀, as an air quality parameter of Karatay, Konya sampling station from beginning January of 2018 to middle of June 2020 (Figure 9) which shows that PM₁₀ values decreased during a pandemic outbreak. PM₁₀ values during the summer hot period lower than colder winter periods.




Figure 9. Daily mean values of PM₁₀, as an air quality parameter of Karatay, Konya sampling station from the end of March of 2018 to middle of October 2019.

Daily mean values of PM₁₀, as an air quality parameter of Selcuklu municipality, Konya sampling station from beginning January of 2018 to middle of June 2020 (Figure 9) which shows that PM₁₀ values decreased during a pandemic outbreak. PM₁₀ values during the summer hot period lower than colder winter periods. Eight month values from middle of December 2018 are missing for some problem of the sampling equipment in the station.




Figure 10. Daily mean values of PM₁₀, as an air quality parameter of Selcuklu municipality, Konya sampling station from beginning January of 2018 to middle of June 2020.

Daily mean values of PM₁₀, as an air quality parameter of Erenkoy, Konya sampling station from end July of 2018 to end of December 2018 (Figure 9) PM₁₀ values were higher during colder winter periods. Other values than 4 months are missing for some problem of the sampling equipment in the station.




Figure 11. Daily mean values of PM₁₀, as an air quality parameter of Erenkoy, Konya sampling station from beginning January of 2018 to middle of June 2020.

Figure 12 explains the monthly mean of PM₁₀ levels for three different sampling stations at Meram, Karatay, and Selcuklu districts of Konya city center from beginning January of 2018 to middle of June 2020. When data were being examined the monthly mean of PM₁₀ values for three different sampling stations shows that winter periods are high and summer periods are lower in a comparison between seasonal rates. Solid lines show the period of March 16 to end of May which is March 16 to end of May 2020 is significantly lower than 2018 and 2019 periods. Table 1 is summarizing the statistical reduction during period of pandemic outbreak time (16 March to end of April 2020) which is lower than before quarantine period (1 January to 15 March 2020) and also lower than similar time scale of 2018 and 2019 period. Monthly mean of PM_{2.5} values during pandemic outbreak quarantine period (01 January to 15 March 2020) were also lower than before quarantine period (1 January to 15 March 2020) and same period of 2018 and 2019.




Figure 12. Monthly mean of PM₁₀ values for three different sampling stations at Karatay and Selcuklu districts of Konya city center from beginning January of 2018 to middle of June 2020.

PM, air quality parameters of Konya all sampling station which they show PM₁₀ and PM_{2.5}, values decreased during COVID-19 pandemic outbreak. Some values of sampling stations are missing and some calculations are not possible (n.a.). It is clearly has been shown that during pandemic quarantine period, air quality was improved (Table 1).

Table 1. Decrease of PM₁₀ and 2.5 micrometer particle size values during pandemic outbreak quarantine period (1 January to 15 March 2020)

	PM ₁₀ (µg/m ³)			PM _{2.5} (µg/m ³)		
	2018	2019	2020	2018	2019	2020
Period 1	72	116	42	n.a.	n.a.	35
Period 2	116	35	25	n.a.	26	22
Period 2/ Period 1	61	-70	-40	n.a.	n.a.	-36
Term2 (2020) - Mean Term2 (2018, 2019)			-66			n.a.

4. CONCLUSION

Due to the continental climate of the Konya province, the winter season is very cold. Therefore, air pollution emissions to the atmosphere increase depending on the amount of fossil fuel used for heating system of heating of residences and other buildings. Meteorological conditions are also important airborne periods determinant in air pollution, since emission are important in air pollution. Covidien-19 coincides the emergence of the pandemic in Turkey that period that corresponds with the end of the winter, although air pollution is also reduced with increasing temperature. The restrictions imposed by the pandemic, besides decreasing the traffic density, caused the emissions to decrease as it also reduces the industry activities. Therefore, the decrease in the level of air pollutants between before and after the restriction is more pronounced compared to previous years. Thus, air quality of restrictions applied during the Covid-19 pandemic quarantine measures greatly influenced positively. Especially the improvement in air quality may be considered as one of important gain with the pandemic measures. Air quality monitoring studies are a great importance in terms of determining the causes and sources of pollution. Their distribution, developing appropriate control strategies, and controlling the effectiveness of these strategies. It was revealed that the measures are taken in the period of a virus outbreak, prohibition of traffic and the reduction of industrial activities caused, a remarkable improvement in air quality. Considering the measures and improvements in air quality, it is considered to be worth investigating how the measures should be evaluated in the coming periods in combating air pollution.

Future Recommendations

Air pollution and Environment affect each other through complex interactions in the atmosphere depending by meteorological factors. Covid-19 pandemic quarantine measures should be taken to

improve air quality and air quality standards should be well implemented. Depending on the ecological change, the related with air pollution should be well emphasized and more discussed in the situation of epidemic period. To increase the sustainable environment and air quality, gains from the measures should be maintained. environmental quality has decreased, as in many parts of the world with the end of the measures.

Acknowledgements: The authors are grateful to the Ministry of Environment and Urbanism of Turkey, Turkish State Meteorological Service for air pollution and meteorological data

REFERENCES

- Arkin, F, (2020) *Asian COVID-19 lockdowns clear the air of pollutants*. Sci. Dev. Net. <https://www.scidev.net/asia-pacific/environment/news/asian-covid-19-lockdownsclear-the-air-of-pollutants.html>.
- Bao, R, Zhang A, (2020). Does lockdown reduce air pollution? Evidence from 44 cities in northern China. *Sci. Total Environ.* **731**. 139052.10.1016/j.scitotenv.2020.139052.
- Chen H, Guo J, Wang C, Luo F, Yu X, Zhang W, Li J, Zhao D, Xu D, Gong Q, Liao J, Yang H, Hou W, Zhang Y, (2020) Clinical characteristics and intrauterine vertical transmission potential of COVID-19 infection in nine pregnant women: a retrospective review of medical records. *Lancet* [https://doi.org/10.1016/S0140-6736\(20\)30360-3](https://doi.org/10.1016/S0140-6736(20)30360-3).
- CNN, (2020) doi. <https://edition.cnn.com/2020/04/09/politics/airline-passengers-decline/index.html>.
- Dutheil F, Baker SJ, Navel V, (2020) COVID-19 as a factor influencing air pollution? *Environ.Pollut.* <https://doi.org/10.1016/j.envpol.2020.114466>.
- Huang, C, Wang, Y, Li, X, Ren, L, Zhao, J, Hu, Y, Zhang, L, Fan, G, Xu, J, Gu, X, Cheng, Z, Yu, T, Xia, J, Wei, Y, Wu, W, Xie, X, Yin, W, Li, H, Liu, M, Xiao, Y, Gao, H, Guo, L, Xie, J, Wang, G, Jiang, R, Gao, Z, Jin, Q, Wang, J, Cao, B, 2020. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet* 395, 497–506. [https://doi.org/10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5).
- Kanniah K, Zaman K, Nurul, Kaskaoutis D, Latif MT, (2020). COVID-19's impact on the atmospheric environment in the Southeast Asia region. *Science of The Total Environment.* **736**. 139658. 10.1016/j.scitotenv.2020.139658.
- Khan MF, Hamid AH, Bari MA, Tajudin AB, Latif MT, 2019. Airborne particles in the city center of Kuala Lumpur: Origin, potential driving factors, and deposition flux in human respiratory airways, *Science of The Total Environment* 650, Part 1, 1195-1206.
- Muhammad S, LongX, (2020). COVID-19 pandemic and environmental pollution: A blessing in disguise? *Sci. Total Environ.* **728**. 138820. 10.1016/j.scitotenv.2020.138820.
- Otmani A, Benchrif A, Mounia T, Bounakhla M, Chakir, E, Bouch M, Krombi M, (2020). Impact of Covid-19 lockdown on PM₁₀, SO₂ and NO₂ concentrations in Salé City (Morocco). *Sci. Total Environ.* **735**. 139541. 10.1016/j.scitotenv.2020.139541.
- URL-1 <http://webgisCovid19.beyond-eocenter.eu/index.php> <http://webgisCovid19.beyond-eocenter.eu/index.php>
- URL-2 <https://www.carbonbrief.org>
- URL-3 <https://www.mgm.gov.tr/veridegerlendirme/il-ve-ilceler-istatistik.aspx?m=KONYA>
- URL-4 <https://www.google.com.tr/maps>
- URL-5 <https://csb.gov.tr/>
- Wang, P, Chen, K, Zhu, S, Wang, P, Zhang, H, 2020. Severe air pollution events not avoided by reduced anthropogenic activities during COVID-19 outbreak. *Resour. Conserv. Recycl.* 158, 104814. <https://doi.org/10.1016/j.resconrec.2020.104814>.
- Zoran M, R.S. Savastru, D.M. Savastru, and N. Marina, Assessing the relationship between surface levels of PM_{2.5} and PM₁₀ particulate matter impact on COVID-19 in Milan, Italy, *Sci. Total Environ.* (2020), <https://doi.org/10.1016/j.scitotenv.2020.139825>
- Zoran M, Savastru R, Savastru D, Penache MC, (2019) Temporal trends of carbon monoxide (CO) and radon (²²²Rn) tracers of urban air pollution. *J. Radio. & Nuclear Chem.* **320**, 5.