

COMPUTATIONAL INVESTIGATION OF THERMOCAPILLARY MIGRATION OF BUBBLES AND DROPS IN ZERO GRAVITY

by
Selman Nas

A dissertation submitted in partial fulfillment
of the requirements for the degree of
Doctor of Philosophy
(Aerospace Engineering)
in The University of Michigan
1995

Doctoral Committee:

Associate Professor Grétar Tryggvason, Chairperson
Associate Professor Luis P. Bernal
Assistant Professor Steve L. Ceccio
Professor Gerard M. Faeth
David Jacqmin, Research Scientist, NASA Lewis Research Center

© Selman Nas 1995
All Rights Reserved

TABLE OF CONTENTS

DEDICATION	ii
ACKNOWLEDGEMENTS	iii
LIST OF FIGURES	vi
CHAPTER	
I. INTRODUCTION	1
1.1 Motivation	1
1.2 Historical Background	2
1.3 Current Work	9
II. FORMULATION AND NUMERICAL METHOD	12
2.1 Governing Equations	12
2.2 Non-dimensional Numbers	14
2.3 Numerical Method	17
2.4 Validation	20
III. THERMOCAPILLARY INTERACTION OF TWO BUB- BLES OR DROPS	32
3.1 Introduction	32
3.2 Resolution test	32
3.3 Thermocapillary migration of a single bubble	33
3.4 Effect of initial position on the interaction of two bubbles	37
3.5 Two dimensional simulations of two bubbles or drops	38
3.6 Three-dimensional simulations of two bubbles	47
3.7 Conclusion	50
IV. THERMOCAPILLARY MIGRATION OF A BUBBLE CLOUD	90
4.1 Introduction	90

4.2	Mono-dispersed systems	90
4.3	Polydispersed systems	94
4.4	Conclusion	99
V. CONCLUSIONS		131
5.1	Conclusions	131
5.2	Suggestions for future work	133
APPENDIX		135
BIBLIOGRAPHY		139

LIST OF FIGURES

Figure

2.1	The computational setup. The top and bottom wall are no-slip boundaries with constant temperature T_{hot} and T_{cold} . The horizontal boundaries are periodic	25
2.2	Velocity versus time. Convergent test for a single two-dimensional bubble on a computational domain which is 4 bubble radii in each direction.	26
2.3	Velocity versus box size for a single bubble in 3d. The resolution is 48^3 grid points, 64^3 grid points and $80^2 \times 96$ grid points, respectively.	26
2.4	The temperature field in the middle plane of the computational domain. The domain is four bubble radii in each direction. The bottom wall is cold and the top wall is hot.	27
2.5	Temperature contours in the middle plane of the computational domain. 50 equally spaced contours are shown.	27
2.6	The velocity field in the middle plane of the computational domain.	28
2.7	Details of the velocity field in the middle plane of the computational domain.	28
2.8	Effect of the viscosity ratio on the migration velocity.	29
2.9	Effect of the conductivity ratio on the migration velocity.	29
2.10	Effect of the density ratio on the migration velocity.	30
2.11	Effect of the heat capacity ratio on the migration velocity.	30
2.12	Schematic of the third test problem.	31

2.13	Resolution test for the problem in Figure 2.12. Total mass flux scaled by the exact mass flux versus grid resolution.	31
3.1	Velocity field (top), isotherms (middle) and streamlines (bottom) of a single bubble rising at steady state. 32×64 grid points of the velocity field, 50 equally spaced isotherms and 21 equally spaced streamlines in a frame moving with the bubble are shown. First column is a 32×64 grid, second column is for a 64×128 grid and third column is for a 128×256 grid.	52
3.2	Resolution test for a single bubble. (a) Migration velocity versus time (b) z component of the trajectories versus time.	53
3.3	Isotherms (top) and streamlines (bottom) of a single bubble migrating at steady state for different Marangoni number. 50 equally spaced isotherms and 20 equally spaced streamlines are shown. Marangoni number for each column from left to right is 1, 10, 20, 40, 60. Here, $Re = 1$ and $Ca = 0.06666$	54
3.4	Effect of Marangoni number on the migration velocity and the deformation of a single bubble.(a) Migration velocity versus time (b) Deformation versus time.	55
3.5	Isotherms (top) and streamlines (bottom) of a single bubble migrating at steady state for different Reynolds number. 50 equally spaced isotherms and 20 equally spaced streamlines are shown. Reynolds number for each column from left to right is 1, 5, 10. Here, $Ma = 1$ and $Ca = 0.06666$ and the resolution is 64×128 grid points.	56
3.6	Effect of Reynolds number on the migration velocity and the deformation of a single bubble. (a) Migration velocity (b) Deformation versus time. Here, $Ma = 1$ and $Ca = 0.06666$	57
3.7	Effect of Capillary number on the migration velocity and the deformation of a single bubble. (a) Migration velocity (b) Deformation versus time. Here, $Re = Ma = 1$	58
3.8	(a) The effect of the viscosity ratio and (b) the effect of the conductivity ratio on the migration velocity of a bubble. Here, $Re = Ma = 1$ and $Ca = 0.06666$	59
3.9	(a) The effect of the density ratio and (b) the effect of the heat capacity ratio on the migration velocity of a single bubble. Here, $Re = Ma = 1$ and $Ca = 0.06666$	60

3.10	6 different initial position for two-bubble interaction. Computational domain size is $x/a = 8$ and $z/a = 16$	61
3.11	The scaled (a) vertical (b) horizontal separation between the bubble on the right and the left versus time. The distance is scaled by bubble radius, a and time is scaled by a/U_r	62
3.12	The scaled migration velocity difference between the bubble on the right and the left versus time. The velocity difference is scaled by U_r and time is scaled by a/U_r	63
3.13	Velocity field for selected frames from the computation of two bubble interaction. Only one third of the velocity field is shown. Time progresses from left to right. The nondimensional time, t^* , is equal to 3.8, 11.4, 25.3, 56.9, 69.5, 88.5, 107.5. The nondimensional time is scaled by a/U_r and velocity is scaled by reference velocity, U_r . Computational domain size is $x/a = 8$ and $z/a = 16$	64
3.14	Temperature contours for selected frames from the computation of two bubble interaction. 50 equally spaced contours are shown. Time progresses from left to right. The nondimensional time, t^* , is equal to 3.8, 11.4, 25.3, 56.9, 69.5, 88.5, 107.5. The nondimensional time is scaled by a/U_r and temperature is scaled, after subtracting a reference temperature, by $a\nabla T_\infty$. Computational domain size is $x/a = 8$ and $z/a = 16$. Here, $Re = Ma = 40$ and $Ca = 0.04166$	65
3.15	Quantitative description of the two bubble interaction in Figure 3.13 and 3.14. (a) Migration velocity versus time. (b) Internal circulation versus time. (c) Deformation versus time. (d) Separation distance versus time. Velocity is scaled by U_r , separation distance by a , time by a/U_r , and the internal circulation by $2\pi aU_r$	66
3.16	Trajectories of the center of mass of the bubbles in Figure 3.13 and 3.14. Both axis are scaled by the bubble radius a	67
3.17	Quantitative information about two-bubble interaction for different Reynolds and Marangoni numbers. (a) The vertical separation distance versus time. (b) The horizontal separation distance versus time. (c) The migration velocity difference versus time. Time is scaled by a/U_r , separation difference, Δ , by a , and the velocity difference by U_r . The differences are between the bubble on the right and the left.	68

3.18	Velocity field for selected frames from the computation of two drop interaction. The velocity field is shown at every third grid point. Time progresses from left to right. The nondimensional time, t^* , is equal to 12.65, 50.60, 75.89, 101.19, 126.49, 151.79, 170.0. The nondimensional time is scaled by a/U_r and velocity is scaled by reference velocity, U_r . Computational domain size is $x/a = 8$ and $z/a = 16$. Here, $Re = Ma = 40$ and $Ca = 0.04166$	69
3.19	Temperature contours for selected frames from the computation of two drop interaction in Figure 3.18. 50 equally spaced contours are shown. Time progresses from left to right. The nondimensional time, t^* , is equal to 12.65, 50.60, 75.89, 101.19, 126.49, 151.79, 170.0. The nondimensional time is scaled by a/U_r and temperature is scaled, after subtracting a reference temperature, by $a\nabla T_\infty$. Computational domain size is $x/a = 8$ and $z/a = 16$	70
3.20	Quantitative description of the two drop interaction in Figures 3.18 and 3.19. (a) Migration velocity versus time. (b) Internal circulation versus time. (c) Deformation versus time. (d) Separation distance versus time. Velocity is scaled by U_r , separation distance by a , time by a/U_r , and the internal circulation by $2\pi aU_r$	71
3.21	Trajectories of the center of mass of drops in Figures 3.18 and 3.19. Both axis are scaled by the drop radius a	72
3.22	Quantitative information for two-drop interactions for different Reynolds and Marangoni numbers. (a) The vertical separation distance versus time. (b) The horizontal separation distance versus time. Time is scaled by a/U_r , separation difference, Δ , by, a . The differences are between the drop on the right and drop on the left.	73
3.23	Velocity field for selected frames from the computation of two bubble interaction. The velocity field is shown at every third grid point. Time progresses from left to right. The nondimensional time, t^* , is equal to 4, 20, 40, 80, 100, 120. The nondimensional time is scaled by a/U_r and velocity is scaled by reference velocity, U_r . Computational domain size is $x/a = 8$ and $z/a = 16$. Here, $Re = Ma = 40$ and $Ca = 0.04166$	74

3.24	Temperature contours for selected frames from the computation of two bubble interaction in Figure 3.23. 50 equally spaced contours are shown. Time progresses from left to right. The nondimensional time, t^* , is equal to 4, 20, 40, 80, 100, 120. The nondimensional time is scaled by a/U_r and temperature is scaled, after subtracting a reference temperature, by $a\nabla T_\infty$. Computational domain size is $x/a = 8$ and $z/a = 16$	75
3.25	Quantitative description of the two bubble interaction in Figures 3.23 and 3.24. (a) Migration velocity versus time. (b) Internal circulation versus time. (c) Deformation versus time. (d) Separation distance versus time. Velocity is scaled by U_r , separation distance by a , time by a/U_r , and the internal circulation by $2\pi aU_r$	76
3.26	Trajectories of the center of mass of the bubbles in Figures 3.23 and 3.24. Both axis are scaled by the bubble radius a	77
3.27	Quantitative information for two-bubble interactions for different Reynolds and Marangoni numbers. (a) The vertical separation distance versus time. (b) The horizontal separation distance versus time. Time is scaled by a/U_r , separation difference, D , by, a . The differences are between the bubble on the right and the left.	78
3.28	Velocity and temperature field for selected frames from the fully three-dimensional simulation of two-bubble interaction. The velocity field is shown at every other grid point in the middle plane of the computational box in y direction. Time progresses from left to right, top to bottom. The nondimensional time, t^* , is equal to 0.57, 5.71, 22.85, 57.14, 85.71. The nondimensional time is scaled by a/U_r and velocity is scaled by reference velocity, U_r . Computational domain size is $x/a = 5.71$, $y/a = 2.86$ and $z/a = 11.43$. Here, $Re = 20$, $Ma = 60$ and $Ca = 0.04166$	79
3.29	Velocity field for selected frames from the fully three dimensional computation of two bubble interaction in Figure 3.28. The velocity field is shown at every other grid point in the middle plane of the computational box in y direction. Time progresses from left to right. The nondimensional time, t^* , is equal to 0.57, 5.71, 22.85, 57.14, 85.71. The nondimensional time is scaled by a/U_r and velocity by U_r . Computational domain size is $x/a = 5.71$, $y/a = 2.86$ and $z/a = 11.43$	80

3.30	Temperature contours for selected frames from the fully three dimensional computation of two bubble interaction in Figure 3.28. 50 equally spaced contours are shown in the middle plane of the computational box in y direction. Time progresses from left to right. The nondimensional time, t^* , is equal to 0.57, 5.71, 22.85, 57.14, 85.71. The nondimensional time is scaled by a/U_r and temperature is scaled, after subtracting a reference temperature, by $a\nabla T_\infty$. Computational domain size is $x/a = 5.71$, $y/a = 2.86$ and $z/a = 11.43$	81
3.31	Quantitative description of the two-bubble interaction in Figures 3.28, 3.29 and 3.30. (a) Migration velocity versus time. (b) z component of the centroid of bubbles versus time. (c) The separation distance versus time and (d) The velocity difference versus time between the bubble on the right and the left. Velocity is scaled by reference velocity U_r , separation distance and z axis by a , and time by a/U_r	82
3.32	Trajectories of the center of mass of the bubbles. Both axis are scaled by the bubble radius a	83
3.33	Velocity and temperature field for selected frames from the fully three-dimensional simulation of two-bubble interaction. The velocity field is shown at every other grid point in the middle plane of the computational box in y direction. Time progresses from left to right, top to bottom. The nondimensional time, t^* , is equal to 3.5, 7.0, 14.0, 35.0, 52.5, 63.0. The nondimensional time is scaled by a/U_r and velocity is scaled by reference velocity, U_r . Computational domain size is $x/a = 5.71$, $y/a = 2.86$ and $z/a = 11.43$. Here, $Re = 60$, $Ma = 20$ and $Ca = 0.04166$	84
3.34	Velocity field for selected frames from the fully three dimensional computation of two bubble interaction. The velocity field is shown at every other grid point in the middle plane of the computational box in y direction. Time progresses from left to right. The nondimensional time, t^* , is equal to 3.5, 7.0, 14.0, 35.0, 52.5, 63.0. The nondimensional time is scaled by a/U_r and velocity by U_r . Computational domain size is $x/a = 5.71$, $y/a = 2.86$ and $z/a = 11.43$	85

3.35	Temperature contours for selected frames from the fully three dimensional computation of two bubble interaction. 50 equally spaced contours are shown in the middle plane of the computational box in y direction. Time progresses from left to right. The nondimensional time, t^* , is equal to 3.5, 7.0, 14.0, 35.0, 52.5, 63.0. The nondimensional time is scaled by a/U_r and temperature is scaled, after subtracting a reference temperature, by $a\nabla T_\infty$. Computational domain size is $x/a = 5.71$, $y/a = 2.86$ and $z/a = 11.43$	86
3.36	Quantitative description of the two-bubble interaction in Figures 3.33, 3.34 and 3.35. (a) Migration velocity versus time. (b) z component of the centroid of bubbles versus time. (c) The separation distance versus time and (d) The velocity difference versus time between the bubble on the right and the left. Velocity is scaled by reference velocity U_r , separation distance and z axis by a , and time by a/U_r	87
3.37	Trajectories of the center of mass of the bubbles. Both axis are scaled by the bubble radius a	88
3.38	Quantitative information for two-bubble interaction for different Reynolds and Marangoni numbers. (a) The vertical separation distance versus time. (b) The horizontal separation distance versus time. Time is scaled by a/U_r , separation difference, Δ , by, a . The differences are between the bubble on the right and the bubble on the left.	89
4.1	Velocity field for selected frames from the computation of six bubble interaction. The velocity field is shown at every third grid point. Time progresses from left to right. The nondimensional time, t^* , is equal to 0.25, 5, 15, 25, 40, 62.5, 87.5, 112.5. The nondimensional time is scaled by a/U_r and velocity is scaled by reference velocity, U_r . Computational domain size is $x/a = 10$ and $z/a = 20$. Here, $Re = 5$, $Ma = 20$ and $Ca = 0.01666$	100
4.2	Temperature contours for selected frames from the computation of six bubble interaction in Figure 4.2. 50 equally spaced contours are shown. Time progresses from left to right. The nondimensional time, t^* , is equal to 0.25, 5, 15, 25, 40, 62.5, 87.5, 112.5. The nondimensional time is scaled by a/U_r and temperature is scaled, after subtracting a reference temperature, by $a\nabla T_\infty$. Computational domain size is $x/a = 10$ and $z/a = 20$	101

4.3	Quantitative description of the six bubble interaction in Figures 4.2 and 4.3. (a) Migration velocity versus time. (b) z component of the centroid of bubbles versus time. (c) Internal circulation versus time. (d) Deformation versus time. Velocity is scaled by U_r , z axis by a , time by a/U_r , and the internal circulation by $2\pi aU_r$. Bubbles marked 1 to 6 from left to right in the domain. Com is the velocity of the center of mass.	102
4.4	Trajectories of the center of mass of the bubbles. Both axis are scaled by the bubble radius a	103
4.5	(a) The average temperature (b) The average temperature fluctuation, versus vertical direction at $t^* = 112.5$. Note that the initial temperature distribution at $t^* = 0.0$ is shown by straight line. Temperature is scaled, after subtracting a reference temperature, by $a\nabla T_\infty$	104
4.6	Velocity field for selected frames from the simulation of 16 bubble interaction. The velocity field is shown at every third grid point. Time progresses from left to right, top to bottom. The nondimensional time, t^* , is equal to 1.4, 14.3, 64.3, 107, 142.85, 214.3, 285.7, 357.14, 428.6. The nondimensional time is scaled by a/U_r and velocity by U_r . Computational domain size is $x/a = 11.43$ and $z/a = 11.43$. Here, $Re = 5$, $Ma = 40$ and $Ca = 0.01666$	105
4.7	Temperature contours for selected frames from the simulation of 16 bubble interaction in Figure 4.6. 50 equally spaced contours are shown. Time progresses from left to right, top to bottom. The nondimensional time, t^* , is equal to 1.4, 14.3, 64.3, 107, 142.85, 214.3, 285.7, 357.14, 428.6. The nondimensional time is scaled by a/U_r and temperature is scaled, after subtracting a reference temperature, by $a\nabla T_\infty$. Computational domain size is $x/a = 11.43$ and $z/a = 11.43$	106
4.8	(a) Migration velocity versus time (b) z component of the centroid of bubbles versus time for 16 equal-sized bubble interaction in Figures 4.6 and 4.7. Velocity is scaled by reference velocity U_r , time is scaled by a/U_r and z axis by a	107
4.9	Trajectories of the center of mass of the bubbles. Both axis are scaled by the bubble radius a	108

4.10	A fully three-dimensional simulation of the interaction of nine equal sized bubbles. Time progresses from left to right, top to bottom. The nondimensional time, t^* , is equal to 0.66, 6.66, 40, 66.66, 80, 86.66. The nondimensional time is scaled by a/U_r and velocity is scaled by reference velocity, U_r . Computational domain size is $x/a = 6.66$, $y/a = 6.66$ and $z/a = 13.33$	109
4.11	Velocity field for selected frames from the fully three dimensional simulation of nine-bubble interaction. The velocity field is shown at every other grid point in the middle plane of the computational box in y direction. Time progresses from left to right. The nondimensional time, t^* , is equal to 0.66, 6.66, 40, 66.66, 80, 86.66. The nondimensional time is scaled by a/U_r and velocity is scaled by reference velocity, U_r . Computational domain size is $x/a = 6.66$, $y/a = 6.66$ and $z/a = 13.33$	110
4.12	Velocity field for selected frames from the fully three dimensional simulation of nine-bubble interaction. The velocity field is shown at every other grid point in the middle plane of the computational box in x direction. Time progresses from left to right. The nondimensional time, t^* , is equal to 0.66, 6.66, 40, 66.66, 80, 86.66. The nondimensional time is scaled by a/U_r and velocity is scaled by reference velocity, U_r . Computational domain size is $x/a = 6.66$, $y/a = 6.66$ and $z/a = 13.33$	111
4.13	Temperature contours for selected frames from the fully three dimensional simulation of nine-bubble interaction. 50 equally spaced contours are shown in the middle plane of the computational box in y direction. Time progresses from left to right. The nondimensional time, t^* , is equal to 0.66, 6.66, 40, 66.66, 80, 86.66. The nondimensional time is scaled by a/U_r and temperature is scaled, after subtracting a reference temperature, by $a\nabla T_\infty$. Computational domain size is $x/a = 6.665$ $y/a = 6.66$ and $z/a = 13.33$	112
4.14	Temperature contours for selected frames from the fully three dimensional simulation of nine-bubble interaction. 50 equally spaced contours are shown in the middle plane of the computational box in x direction. Time progresses from left to right. The nondimensional time, t^* , is equal to 0.66, 6.66, 40, 66.66, 80, 86.66. The nondimensional time is scaled by a/U_r and temperature is scaled, after subtracting a reference temperature, by $a\nabla T_\infty$. Computational domain size is $x/a = 6.66$, $y/a = 6.66$ and $z/a = 13.33$	113

4.15	(a) Migration velocity versus time (b) z component of the centroid of bubbles versus time, for nine-bubble simulation in Figure 4.10. Velocity is scaled by U_r , time by a/U_r and z axis by a	114
4.16	(a) $x - z$ trajectories of the center of mass of the bubbles (left). (b) $y - z$ trajectories of the center of mass of the bubbles (right). Both axis are scaled by the bubble radius a	115
4.17	Velocity field for selected frames from the computation of six unequal-sized-bubble interaction. Third of the velocity field is shown at every third grid point. Time progresses from left to right. The nondimensional time, t^* , is equal to 2.5, 12.5, 25, 40, 62.5, 87.5, 112.5, 137.5. The nondimensional time is scaled by a_v/U_{rv} and velocity is scaled by reference velocity, U_{rv} . Computational domain size is $x/a_v = 10$ and $z/a_v = 20$	116
4.18	Temperature contours for selected frames from the computation of six unequal-sized-bubble interaction. 50 equally spaced contours are shown. Time progresses from left to right. The nondimensional time, t^* , is equal to 2.5, 12.5, 25, 40, 62.5, 87.5, 112.5, 137.5. The nondimensional time is scaled by a_v/U_{rv} and temperature is scaled, after subtracting a reference temperature, by $a_v \nabla T_\infty$. Computational domain size is $x/a_v = 10$ and $z/a_v = 20$	117
4.19	Quantitative description of the six unequal-sized-bubble interaction in Figures 4.17 and 4.18. (a) Migration velocity versus time. (b) z component of the centroid of bubbles versus time. (c) Internal circulation versus time. (d) Deformation versus time. Velocity is scaled by U_{rv} , z axis by the averaged bubble radius, a_v , time by a_v/U_{rv} , and the internal circulation by $2\pi a_v U_{rv}$. Bubbles marked 1 to 6 from left to right in the domain. Com is the velocity of the center of mass.	118
4.20	Trajectories of the center of mass of the bubbles. Both axis are scaled by the average bubble radius a_v	119
4.21	Velocity field for selected frames from the simulation of 16 unequal-sized bubble interaction. The velocity field is shown at every third grid point. Time progresses from left to right, top to bottom. The nondimensional time, t^* , is equal to 4, 40.5, 64.7, 89, 113.3, 137.6, 161.9, 182, 202.4. The nondimensional time is scaled by a_v/U_{rv} and velocity by U_{rv} . Computational domain size is $x/a_v = 13.33$ and $z/a_v = 13.33$	120

4.22	Isotherms for selected frames from the simulation of 16 unequal-sized bubble interaction. 50 equally spaced contours are shown. Time progresses from left to right, top to bottom. The nondimensional time, t^* , is equal to 4, 40.5, 64.7, 89, 113.3, 137.6, 161.9, 182, 202.4. The nondimensional time is scaled by a_v/U_{rv} and temperature is scaled, after subtracting a reference temperature, by $a_v \nabla T_\infty$. Computational domain size is $x/a_v = 13.33$ and $z/a_v = 13.33$	121
4.23	(a) Migration velocity versus time (b) z component of the centroid of bubbles versus time for 16 unequal-sized bubble interaction in Figures 4.21 and 4.22. Velocity is scaled by U_{rv} , time by a_v/U_{rv} and z axis by a_v	122
4.24	Trajectories of the center of mass of the bubbles. Both axis are scaled by the average bubble radius a_v	123
4.25	A fully three-dimensional simulation of the interaction of nine unequal-sized bubbles. Time progresses from left to right, top to bottom. The nondimensional time, t^* , is equal to 6.66, 40, 66.66, 80, 100, 114.2. The nondimensional time is scaled by a_v/U_{rv} and velocity is scaled by average reference velocity, U_{rv} . Computational domain size is $x/a_v = 8$, $y/a_v = 8$ and $z/a_v = 16$	124
4.26	Velocity field for selected frames from the fully three dimensional simulation of nine unequal-sized bubble interaction. The velocity field is shown at every other grid point in the middle plane of the computational box in y direction. Time progresses from left to right. The nondimensional time, t^* , is equal to 6.66, 40, 66.66, 80, 100, 114.2. The nondimensional time is scaled by a_v/U_{rv} and velocity is scaled by average reference velocity, U_{rv} . Computational domain size is $x/a_v = 8$, $y/a_v = 8$ and $z/a_v = 16$	125
4.27	Velocity field for selected frames from the fully three dimensional simulation of nine unequal-sized bubble interaction. The velocity field is shown at every other grid point in the middle plane of the computational box in x direction. Time progresses from left to right. The nondimensional time, t^* , is equal to 6.66, 40, 66.66, 80, 100, 114.2. The nondimensional time is scaled by a_v/U_{rv} and velocity is scaled by average reference velocity, U_{rv} . Computational domain size is $x/a_v = 8$, $y/a_v = 8$ and $z/a_v = 16$	126

- 4.28 Temperature contours for selected frames from the fully three dimensional simulation of nine unequal-sized bubble interaction. 50 equally spaced contours are shown in the middle plane of the computational box in y direction. Time progresses from left to right. The nondimensional time, t^* , is equal to 6.66, 40, 66.66, 80, 100, 114.2. The nondimensional time is scaled by a_v/U_{rv} and temperature is scaled, after subtracting a reference temperature, by $a_v \nabla T_\infty$. Computational domain size is $x/a_v = 8$, $y/a_v = 8$ and $z/a_v = 16$. . . 127
- 4.29 Temperature contours for selected frames from the fully three dimensional simulation of nine unequal-sized bubble interaction. 50 equally spaced contours are shown in the middle plane of the computational box in x direction. Time progresses from left to right. The nondimensional time, t^* , is equal to 6.66, 40, 66.66, 80, 100, 114.2. The nondimensional time is scaled by a_v/U_{rv} and temperature is scaled, after subtracting a reference temperature, by $a_v \nabla T_\infty$. Computational domain size is $x/a_v = 8$, $y/a_v = 8$ and $z/a_v = 16$. . . 128
- 4.30 (a) Migration velocity versus time (b) z component of the centroid of bubbles versus time, for 9 unequal-sized-bubble simulation. Velocity is scaled by U_{rv} , time by a_v/U_{rv} and z axis by average bubble radius, a_v . Even numbered bubbles are the larger ones. 129
- 4.31 (a) $x - z$ trajectories of the center of mass of the bubbles (left). (b) $y - z$ trajectories of the center of mass of the bubbles (right). Both axis are scaled by the average bubble radius, a_v 130